

August 31, 2022

Mint Pest Alert Newsletter

-Central Oregon-

Fall Chemical Management

Late Aug/early Sep is the time to scout for Mint Root Borer (MRB) larvae in the soil if fall control is being considered (i.e. Lorsban, Mocap, nematodes). MRB larvae drop from the foliage to the ground in August and tunnel into rhizomes, where they will feed into the fall.

Larvae are generally too small to see in August, while sampling in late September is too late as damage is done.

Control treatments must be applied before larvae form hibernacula (overwintering, soil-based cocoons). Based on the insect development model, hibernacula formation in the Madras area reached 5% hibernacula formation on August 28 and will reach 50% by Sept 12. The Powell Butte area should expect 5% hibernacula by September 6 and 50% by October 7.



Oregon State University
Extension Service

Questions? Contact

John Spring: John.spring@oregonstate.edu

To sign up for this newsletter contact

Christy Tanner: Christy.tanner@oregonstate.edu

Sponsored by the Oregon Mint Commission

OREGONmint



Sampling Recommendations for MRB larvae:

- Sample 1ft² of soil to a depth of 2-3" (include rhizomes)
- 2 samples taken every 2-3 acres
- Minimum 25 samples per field
- Screen samples in the field or with Berlese funnels in the laboratory

Treatment threshold = average of 2-3 larvae/ft² sample

